





| 7 | NCBI | | Pub | W(e | d | | ibrary dicine | | |
|---|---------|---|------------|-------------|--------------|---------|------------------|-------------|----------|
| Entrez | PubMed | Nucleotide | Protein | Genome | Structure | ОМІМ | PMC | Journals | В |
| Search | PubMed | for | | | | Go | Clear | | |
| | | Limits | Previ | ew/Index | History | CI | lipboard | De | etails |
| About En | trez | | | | | | | | |
| | | Display Abs | stract | Sho | ow: 20 ▼ S | Sort | ▼ Ser | nd to Text | <u> </u> |
| Text Vers | ion | | _ | | (a) 10a ana | | | | |
| | 4. 6. 6 | 1: Gastroe | | 1980 Sep;79 | (3):493-502. | | | Related Art | icles, l |
| Entrez PubMed Overview | | Gastroenterology | | | | | | | |
| Help FA | | Immunocytochemical localization of human intrinsic factor: the | | | | | | | |
| Tutorial New/Noteworthy E-Utilities | | nonstimulated stomach. | | | | | | | |
| | | Levine JS, Nakane PK, Allen RH. | | | | | | | |
| PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby Related Resources Order Documents NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central Privacy Policy | | Human intrinsic factor (IF) is found on the surface of parietal cells at the ligh microscopic level by immunofluorescence, but there is no direct evidence the IF is synthesized in and secreted from this cell. In order to provide this evidence, IF was localized at the ultrastructural level in human stomach, usin an immunochemically, monospecific antibody against purified human IF. We defined optimal immunocytochemical conditions and identified the subcellul location of IF in fundic biopsies from 4 fasting subjects. Intrinsic factor was found on the perinuclear membrane, on rough endoplasmic reticulum, Golgi apparatus, tubulovesicles, and membranes of the multivesicular body in paric cells. Intrinsic factor was absent from nuclei, mitochondria, basolateral surfamembranes, blunt microvilli, and the cytosol. Thus, we have defined the paricell as the site of IF synthesis. The observed distribution of intracellular IF suggested that an interrelationship existed between the stained membranous structures, and that IF secretion may depend upon membrane translocation. Further immunocytochemical and physiologic studies should increase our understanding of the cellular physiology of IF secretion as well as the parieta cell. | | | | | | | |
| | | PMID: | 7429110 [F | PubMed - in | dexed for ME | EDLINE] | l | | |

Show: 20 Sort Send to Display Abstract

> Write to the Help Desk NCBI | NLM | NIH
> Department of Health & Human Services Freedom of Information Act | Disclaimer

> > Jan 29 2004 15: